

Vaccines in pregnancy can help protect you, and your unborn child.

Vaccine for pregnant women:

- Around 20 weeks whooping cough (pertussis) vaccine
- From 28 weeks – RSV vaccine
- Flu jab, if pregnant during flu season

Contact your GP if:

- You or your children have missed any vaccinations
- You or your child have vaccination appointments but you've missed it or cannot attend
- Your GP surgery can book or rearrange an appointment

School-aged immunisation service at school

Children and young people can get their vaccines at school.

For more information:



Keep me! NHS vaccination checklist

For the best protection, it's important your child gets their vaccines on time.

Check their Red Book for their vaccination record, and contact your GP to catch up if they've missed any.



1 weeks

- 6-in-1 vaccine (1st dose)
- Rotavirus vaccine (1st dose)
- MenB vaccine (1st dose)

12 weeks

- 6-in-1 vaccine (2nd dose)
- MenB vaccine (2nd dose)
- Rotavirus vaccine (2nd dose)

16 weeks

- 6-in-1 vaccine (3rd dose)
- Pneumococcal vaccine (1st dose)
- Vaccines for children aged one to 15 years

8 year

- MMRV vaccine (1st dose for children born on or after 1 January 2025)
- Pneumococcal vaccine (2nd dose)
- MenB vaccine (3rd dose)

18 months

- 6-in-1 vaccine (4th dose for children born on or after 1 July 2024)

- MMRV vaccine (1st or 2nd dose for children born on or after 1 July 2024)

3 years and 4 months

- 4-in-1 pre-school booster vaccine
- MMRV vaccine (1st or 2nd dose for children born between 1 September 2022 and 31 December 2024)

12 to 13 years

- HPV vaccine

14 years

- Td/IPV vaccine (3-in-1 teenage booster)
- MenACWY vaccine

2 to 15 years

- Children's flu vaccine (every year until children finish Year 11 of secondary school)

More information about childhood vaccines:

